



FASTMath Tech Area Summary: Time Integrators

Carol Woodward (LLNL)



Rensselaer



SMU



USC University of Southern California

The time integrator area is developing time integration methods and software for use in SciDAC applications

Includes two major efforts:

- Multirate methods in the Suite of Nonlinear and Differential-Algebraic Solvers (SUNDIALS)
 - Developing new methods and utilizing some developed in the base program work by Sandu et al.
- Parallel-in-time methods in SUNDIALS utilizing the XBraid software
 - XBraid supported by base program

Synergistic activities between time integration and other FASTMath:

- Spectral deferred correction methods and software for structured grids (in AMReX)
 - New methods planned for AMReX may include hybrid SDC-exponential integrators being developed in base program
- Multirate methods for structured grid systems (SUNDIALS with AMReX)
- Time integrators with adjoint sensitivities for optimization (PETSc)
 - Utilized by MMiCCS center and base program projects



The time integrator team and its synergistic activities span three laboratories and one university

LLNL: Carol Woodward, David Gardner, and John Loffeld

Southern Methodist University: Daniel Reynolds

LBNL: Michael Minion

ANL: Hong Zhang, Barry Smith, Alp Dener

SciDAC application interactions

- (BER) Assessing and Improving the Numerical Solution of Atmospheric Physics in E3SM, PI: Hui Wan (PNNL)
 - Assessing time integration convergence in E3SM climate physics and improving time integration accuracy/stability
- (FES) Partnership for Multiscale Gyrokinetic Turbulence, PI: David Hatch (UT Austin) – See Darin Ernst's talk
 - Interfacing IMEX and multirate methods from SUNDIALS ARKode in 2D test code
 - Interfacing KINSOL nonlinear solver from SUNDIALS to provide acceleration for fixed point solve in Tango code
- (BER) A Non-hydrostatic Variable Resolution Atmospheric Model in ACME, PI: Mark Taylor (SNL)
 - Used ARKode in new E3SM nonhydrostatic dynamical core to test over 100 integration approaches and determine most efficient
 - Ongoing support of ARKode use in E3SM

Through FASTMath we are developing high order multirate time integrators in SUNDIALS

- Applications will become more multiphysics and multirate as they take advantage of exascale systems
- Many commonly used splitting schemes can suffer from low accuracy and/or poor/unknown stability
- Greater flexibility in time integration libraries is needed to meet the needs of multiphysics applications
- We are expanding the capabilities of the SUNDIALS ARKode package to meet these needs
 - Adding adaptive step multirate integration methods

$$y' = f_{\text{fast}}(y) + f_{\text{slow}}(y)$$

We restructured the SUNDIALS ARKode package to provide infrastructure for general, adaptive, one-step methods

- ARKode provides the outer time integration loop and generic usage modes
- Time-stepping modules handle problem-specific components

Released modules:

- *ARKStep* – single-rate explicit, implicit, and IMEX RK methods
- *ERKStep* – a streamlined module for explicit methods
- *MRISep* – module for multirate infinitesimal step methods
 - SUNDIALS 4.0: ARKode MRISep module supporting explicit/explicit 2nd and 3rd order MIS methods.
 - SUNDIALS 5.0-dev.0: Support for implicit and IMEX fast methods and adaptive fast time steps in MRISep

In development: *IMEXGARKStep* – module for IMEX Generalized Additive Runge-Kutta



See poster by David Gardner

1D Advection with Brusselator Reactions

$$\begin{aligned}\frac{\partial u}{\partial t} &= -c \frac{\partial u}{\partial x} + k_1 A - k_2 B u + k_3 u^2 v - k_4 u \\ \frac{\partial v}{\partial t} &= -c \frac{\partial v}{\partial x} + k_2 B u - k_3 u^2 v\end{aligned}$$

- Compare 3rd order IMEX and MIS (multirate) methods
- Explicit (slow) advection and implicit (fast) reactions


	IMEX	MRI
Steps	6,805	1,429 / 8,813
Advection evals	29,379	4,288
Reaction evals	98,917	116,949

Same solution quality
with 85% fewer advection
evaluations (reduces
parallel communication)

- Gains with MRI depend on the separation of time scales e.g., CFL vs reaction rates.

We will be adding a parallel-in-time capability to SUNDIALS through multigrid reduction in time (MGRIT)


- MGRIT uses multiple coarse grids to update the fine grid solution



- Addressing challenges associated with multistep methods and use of variable step adaptive methods
- Initial work (through Off. Electricity project) shows parallel speedups with multi-stage and multistep adaptive step methods
- In year 3 we will connect the LLNL XBraid package with SUNDIALS to deliver MGRIT capability

General support for spectral deferred corrections integration enabled in AMReX structured adaptive mesh framework

- Large-scale computing platforms allow for highly complex multiphysics simulations, but these systems are often integrated with low-order splittings
 - SDC provides high-order time integration through iterations over the native operator split solution procedure
 - Prior support for SDC was specific to certain applications
 - Code and tutorial included in AMReX repo
 - FY18 work provided the capability to allow straightforward construction of high-order semi-implicit and multi-implicit integrators for problems with one or more stiff terms
 - FY19 work extends SDC to multirate integration for AMR



High-order convergence of a multi-implicit SDC integrator run on a 2D advection-diffusion-reaction system

We are collaborating with the structured mesh team to investigate three-rate IMEX methods for AMR grids

- Working with the AMReX team, we are developing multirate methods for advection-diffusion-reaction systems

$$\begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = f_a(y_1) + f_d(y_1) + f_r(y_2)$$

- Methods will include both implicit and explicit algorithms
- Implementation will be in ARKode and tested through an interface within AMReX

We have enabled some multirate time stepping of AMReX problems using SUNDIALS


Currently parallel single-level and multi-level time stepping if Δt is the same on all levels:

- SUNDIALS wrap of AMReX vector structure
- IMEX, or implicit stepping if the implicit portion is in the form $(\alpha A - \beta \nabla \cdot B \nabla) \phi = f$
 - Wrap of AMReX La Placian linear solver as a SUNDIALS linear solver package
 - Two-rate implicit-fast methods implemented. Implicit-slow methods in progress
- Demonstrated via multirate stepping of single- and multi-level Gray-Scott (two-rate) reaction-diffusion problem


See poster by John Loffeld

Gray-Scott equation computed with third-order explicit-explicit MIS multirate method

$$\frac{\partial u}{\partial t} = A \nabla^2 u - uv^2 + f(1 - u), \quad \frac{\partial v}{\partial t} = B \nabla^2 v + uv^2 - (f + k)v$$





$t = 1000$




$t = 5000$

Sensitivity analysis integrator tools for PDE-constrained optimization

- **New composable and scalable discrete-adjoint solvers in PETSc for local sensitivity analysis**
 - Used discretize-then-differentiate approach, overcomes difficulties with Automatic Differentiation
 - Optimal adjoint checkpointing utilizes hierarchical memory
 - Implementation allows multiple objective functions and time intervals
- **Efficient gradients and Hessian-vec products for PDE-constrained opt**
 - Applications in climate data assimilation and power grid
 - New 2nd-order adjoints accelerate convergence of optimization algs



Adjoint-based
4D-Var global
data assimilation



Optimal economic dispatch for dynamic power system

S. Abhyankar, J. Brown, E. Constantinescu, D. Ghosh, B. Smith, H. Zhang, PETSc/TS: A Modern Scalable ODE/DAE Solver Library, arXiv:1806.01437

Our work in ECP provides greater capabilities for SciDAC applications



- SUNDIALS project in ECP
 - Redesign of nonlinear and linear solver interfaces
 - Encapsulation of nonlinear solvers
 - Addition of many-vector interface
- Participating in math libraries software development kit, PI: Yang; developing interoperability layers with SuperLU_DIST, PETSc, *hypre*, AMReX, MFEM, etc.
- Working with AMReX team to optimize use of SUNDIALS for reaction systems on GPUs

FASTMath integrators are used in numerous non-SciDAC DOE applications

- (OE) Supporting use of the IDA DAE integrator in the development of the GridDyn time domain power grid simulation package, PI: Philip Top (LLNL)
- (BER) Supporting use of KINSOL nonlinear solvers in the ParFlow watershed simulation package, PI: Reed Maxwell (CO School of Mines)
- (FES) BOUT++ fusion code uses CVODE as its time integrator
- (ECP) Supporting use of CVODE as an option in AMReX as a reaction integrator; through AMReX have impact on the Nyx cosmology code
- (ECP) CVODE used in the NEK5000 code for reactor modeling
- Numerous other applications throughout DOE and worldwide



CASC

Center for Applied
Scientific Computing



**Lawrence Livermore
National Laboratory**

Disclaimer

This document was prepared as an account of work sponsored by an agency of the United States government. Neither the United States government nor Lawrence Livermore National Security, LLC, nor any of their employees makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States government or Lawrence Livermore National Security, LLC. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States government or Lawrence Livermore National Security, LLC, and shall not be used for advertising or product endorsement purposes.